

604405
604405

1

ON A QUASI-LINEAR EQUATION

Richard Bellman

P-575 ✓

Bev

1 October 1954

Approved for OTS release

11 p



COPY	1	OF	1	p
HARD COPY	\$.1.00			
MICROFICHE	\$.650			

DDC
RECEIVED
AUG 27 1964
DDC-IRA D

The RAND Corporation

1700 MAIN ST • SANTA MONICA • CALIFORNIA

SUMMARY


 The purpose of this paper is to establish some limit
 theorems for the solutions of ~~$x_1(n+1) = \max_{1 \leq j \leq q} a_{1j}(q)x_j(n)$~~ *non-linear recurrence relations.*
 Recurrence relations of this kind occur in various dynamic
 programming problems. () 

i

ON A QUASI-LINEAR EQUATION

Richard Bellman

§1. Introduction

The purpose of this note is to establish some limit theorems for ~~the~~ non-linear recurrence relations

$$(1) \quad x_1(n+1) = \text{Max}_q \sum_{j=1}^n a_{1j}(q)x_j(n), \quad j = 1, 2, \dots, n, \quad n \geq 0,$$

under certain assumptions concerning the initial values $c_1 = x_1(0)$, and the coefficient matrices $A(q) = (a_{1j}(q))$.

Equations of this type occur in various parts of the theory of dynamic programming, as we shall indicate below, and are, in addition, of interest in furnishing a link between the theory of linear and non-linear operations, as we have discussed elsewhere, cf [1], [2], [3], [4].

§2. The Homogeneous Equation

Let us consider the equation

$$(1) \quad \lambda y_1 = \text{Max}_q \sum_{j=1}^n a_{1j}(q)y_j, \quad i = 1, 2, \dots, n,$$

where we impose the following conditions.

- (2) (a) $q = (q_1, q_2, \dots, q_n)$ runs over some set of values, S , with the property that the maximum is attained in (1),
- (b) $\infty > m \geq a_{1j}(q) > 0$, $i, j = 1, 2, \dots, n$ for $q \in S$,
- (c) for any q , let $\phi(q)$ denote the characteristic root of $A(q) = (a_{1j}(q))$ of largest absolute value, the Perron

root, known to be positive. We assume that there exists at least one value of q for which $\phi(q)$ assumes its maximum for $q \in S$.

We shall now prove

Theorem 1. Under these conditions, there exists a unique positive λ with the property that (1) has a positive solution, $y_i > 0$, $i=1,2,\dots,n$. This solution is unique up to a multiplicative constant, and

$$(3) \quad \lambda = \max_{q \in S} \phi(q).$$

Proof. We begin by showing the existence of a positive λ and a positive set of solutions $\{y_i\}$. Consider the region defined by $y_i \geq 0$, $\sum_{i=1}^n y_i = 1$. The normalized transformation

$$(4) \quad y_i' = \left[\max_q \sum_{j=1}^n a_{ij}(q) y_j \right] / \left[\sum_{i=1}^n \max_q \sum_{j=1}^n a_{ij}(q) y_j \right],$$

is a continuous mapping of this region into itself. Hence there exists a fixed point, $\{y_i\}$. This fixed point is a solution of (1), with λ the denominator in (4). Each component y_i is positive because of the positivity of $a_{ij}(q)$.

To show that this solution is unique up to a multiplicative constant, let $[\mu, z]$ be another solution of (1) with $\mu > 0$ and z a positive vector. Let $\{q\}$ be the set of values for which the maximum is attained in (1), and $\{\bar{q}\}$ the similar set associated with z . Observe that we may have different sets for each i . We have then

$$(5) \quad \lambda y_1 = \sum_j a_{1j}(q)y_j \geq \sum a_{1j}(\bar{q})y_j, \quad 1 = 1, 2, \dots, n,$$

$$\mu z_1 = \sum_j a_{1j}(\bar{q})z_j.$$

Let us now assume, without loss of generality that $\lambda < \mu$. Let ϵ be a positive constant chosen so that one, at least, of the components $y_1 - \epsilon z_1$ is zero, one at least is positive, and the others are non-negative. This can always be accomplished if y and z are not proportional. If 1 is an index for which $y_1 - \epsilon z_1$ is zero, we have

$$(6) \quad 0 = \mu(y_1 - \epsilon z_1) > \lambda y_1 - \epsilon \mu z_1 \geq \sum_{j=1}^n a_{1j}(\bar{q})(y_j - \epsilon z_j) > 0,$$

Since $a_{1j}(\bar{q}) > 0$, a contradiction. Hence $\lambda = \mu$ and y and z are proportional.

To show that $\lambda = \text{Max}_q \phi(q)$, we proceed as follows. Let $\mu = \text{Max}_q \phi(q)$. It is clear that λ , as the characteristic root of some $A^q(q)$, satisfies the inequality $\lambda \leq \mu$. Assume that actually $\mu > \lambda$. Let $z = (z_1, z_2, \dots, z_n)$ be a positive characteristic vector associated with μ and \bar{q} a set of q -values which yield $\mu = \phi(\bar{q})$. Then we have

$$(7) \quad \mu z_1 = \sum_{j=1}^n a_{1j}(\bar{q})z_j \leq \text{Max}_q \sum_{j=1}^n a_{1j}(q)z_j$$

Since y_1 is positive, we can find a positive constant m such that $z_1 \leq m y_1$ for $1 = 1, 2, \dots, n$. Hence, (7) yields

$$(8) \quad \mu z_1 \leq m \text{Max}_q \sum_{j=1}^n a_{1j}(q)y_j = m \lambda y_1$$

Thus $z_1 \leq my_1 \lambda/\mu$. Iterating this we obtain $z_1 \leq my_1 (\lambda/\mu)^k$, for arbitrary k . Since $\lambda/\mu < 1$, by assumption, this yields $z_1 = 0$, a contradiction. Hence $\lambda = \mu$.

§3. The Recurrence Relation

Let us now return to the recurrence relation of (1.1) and prove

Theorem 2. If, in addition to the conditions of (2.2), we assume that there is a unique q for which the maximum value of $\phi(q)$ is attained and that $c_1 > 0$, then

$$(1) \quad x_1(n) \sim ay_1 \lambda^n,$$

as $n \rightarrow \infty$, where a is a constant dependent upon the initial values c_1 .

Proof. Let us take $c_1 > 0$ without loss of generality. There are then two positive constants k and K such that $ky_1 \leq c_1 \leq Ky_1, i=1,2,\dots,n$.

Let us show inductively that

$$(2) \quad ky_1 \lambda^n \leq x_1(n) \leq Ky_1 \lambda^n.$$

Assume that we have the result for n , then

$$(3) \quad \begin{aligned} x_1(n+1) &\leq K\lambda^n \max_q \sum_{j=1}^n a_{1j}(q)y_j = K\lambda^{n+1}y_1 \\ &\geq k\lambda^n \max_q \sum_{j=1}^n a_{1j}(q)y_j = k\lambda^{n+1}y_1. \end{aligned}$$

To establish the asymptotic behavior we show that for n sufficiently large the set of q 's which furnish the maximum in (1.1) is precisely the set which yields $\lambda = \max \phi(q)$.

Assume the contrary. This means that infinitely often we employ a set $\{\bar{q}\}$ which is not identical with the q which furnishes the maximum in $\phi(q)$.

We then have

$$(4) \quad x_1(n+1) = \sum_{j=1}^n a_{1j}(\bar{q})x_j(n), \quad i=1,2,\dots,n \\ \leq \left(\sum_{j=1}^n a_{1j}(\bar{q})y_j\right) K\lambda^n.$$

For some index i we must have

$$(5) \quad \sum_{j=1}^n a_{1j}(\bar{q})y_j < \lambda y_1,$$

with strict inequality. For if $\sum_{j=1}^n a_{1j}(\bar{q})y_j \geq \lambda y_1$ for all i , the characteristic root of $A(\bar{q}) = (a_{ij}(\bar{q}))$ of largest absolute value, $\phi(\bar{q})$, would at least equal $\lambda = \max_q \phi(q)$, which would contradict the assumption concerning the uniqueness of the maximum of $\phi(q)$.

Hence, for some component, say the first, we have

$$(6) \quad x_1(n+1) \leq \theta K\lambda^{n+1}y_1, \quad 0 < \theta < 1.$$

Since $a_{1j}(\bar{q}) > 0$ for i,j , where \bar{q} is the value of q for which $\lambda = \phi(\bar{q})$, we see that, for $i = 1,2,\dots,n$,

$$(7) \quad x_1(n+2) \leq K\lambda^{n+1} \left[\sum_{j=2}^n a_{1j}(\bar{q})y_1 + \theta a_{11}(\bar{q})y_1 \right] \\ \leq \theta_1 K\lambda^{n+2}y_1,$$

where $\theta < 1$.

If therefore a set of q 's distinct from q^* are used R times, we obtain

$$(8) \quad x_1(n) \leq \Theta_1^R K \lambda^n y_1,$$

for n sufficiently large. Since $0 < \Theta_1 < 1$, if R is too large we eventually contradict the lower bound for $x_1(n)$.

Hence for $n \geq n_0 = n_0(c_1)$, we have

$$(9) \quad x(n+1) = A(q^*) x(n),$$

whence the asymptotic statement of (1) follows.

§4. A Dynamic Programming Problem

Suppose that we are engaged in a multi-stage decision process of the following type. At each stage we have our choice of various operations, which we number $i=1,2,\dots,K$. The i^{th} operation has a probability distribution attached with the following properties:

- (1) a. There is a probability p_{1k} that we receive k units and the process continues, $k=1,2,\dots,R$,
- b. There is a probability p_{10} that we receive nothing and the process terminates.

How do we proceed so as to maximize the probability that we receive at least n units before the process terminates?

Let us define the sequence

- (2) $u(n)$ = the probability of attaining at least n units before the termination of the process using an optimal procedure.

Then using the intuitive "principle of optimality", cf. [1], [2], [3], [4], we see that $u(n)$ satisfies the recurrence relation

$$(3) \quad u(n) = \max_1 \left[\sum_{k=1}^R p_{1k} u(n-k) \right], \quad n > 0$$

$$= 1, \quad n \leq 0.$$

Using methods similar to those above, we see that for large n ,

$$(4) \quad u(n) \sim c \rho^n,$$

where ρ is the root of largest absolute value, necessarily positive, of

$$(5) \quad 1 = \sum_{k=1}^R p_{1k} \rho^{-k},$$

for the value of 1 which maximizes ρ .

§5. An Analogue of a Result of Markoff

Markoff showed that if

$$(1) \quad x_1(n+1) = \sum_{j=1}^n a_{1j} x_j(n), \quad n = 0, 1, \dots,$$

$x_1(0) > 0$, with the conditions

$$(2) \quad a_{1j} > 0, \quad \sum_j a_{1j} = 1, \quad j = 1, 2, \dots, n,$$

then

$$(3) \quad \lim_{n \rightarrow \infty} x_1(n) = c, \quad i = 1, 2, \dots, n,$$

where c depends on the initial values.

The same proof shows that the same result holds for the sequence defined by

$$(4) \quad x_1(u+1) = \max_q \sum_{j=1}^n a_{1j}(q)x_j(n),$$

provided that the conditions in (2) hold uniformly in q . The constant will, of course, in general, be different from that above.

BIBLIOGRAPHY

1. Bellman, R., "An Introduction to the Theory of Dynamic Programming", RAND Report No. R-245, 1953.
2. ———, "The Theory of Dynamic Programming", Bull. Amer. Math. Soc., (to appear).
3. ———, "Some Functional Equations in the Theory of Dynamic Programming", (to appear).
4. ———, "A Survey of the Theory of Dynamic Programming", Chapter XII of Mathematics for Modern Engineers, E. F. Beckenbach, ed., McGraw-Hill, (forthcoming).

md

BLANK PAGE